

ANALYSIS (w/w):

Magnesium Oxide (MgO) -	4.5%
Magnesium (as Mg) -	2.7%
Sulphur Trioxide (SO ₃) -	19.5%
Sulphur (as S) -	7.8%
Iron (Fe) -	4.2%
Manganese (Mn) -	2.0%
Zinc (Zn) -	1.3%

HortiChel Compound Foliar T.E. Fertiliser Complexed with lignosulphonates

HortiChel Compound Foliar T.E. Fertiliser can be applied on its own or tank mixed with HortiStar, HortiCron and HortiPhyte.

Crop	Rate (litres/ha)	Water volume (litres/ha)	Growth stage	Notes
Apples	5.0	200 to 1000	Petal fall	Repeat application at 3 week intervals
Cane fruit newly planted	2.0	200 to 600	From full leaf expansion	Repeat application at 3 week intervals
Cane fruit established	3.0	200 to 1000		
Grapes	3.0	250 to 400		
Hops	3.0	500 to 1000		
Strawberries newly planted	1.0	200 to 600	2 weeks after planting	Repeat application at 3 week intervals
Strawberries established	2.0	200 to 1000	Green bud stage	Repeat application at 3 week intervals
Hardy Nursery Stock	5.0	200 to 600	From start of growth to mid September, monthly	
Pot and Bedding plants	1.0	200 to 600	From early growth stage repeated every 3 weeks	
Trees and amenity plantings	5.0	200 to 1000	From start of growth to mid September, monthly. Apply to leaves and to bark	
Seedlings and cuttings	1.0	400 to 1000	Apply from first true leaf stage to planting at 2 week intervals.	
Turf	5.0	200 to 600	March and September	

Mixing Instructions:

Half fill the spray tank with water. Add the HortiChel Compound T.E. Foliar Fertiliser to the tank and agitate well. Add any other Hortifeeds products and agitate again. Do not store for more than 24 hours once mixed. Apply sprayer washings to the crop. Avoid pollution of watercourses.

HortiChel, HortiStar, HortiCron and HortiPhyte are fertilisers with no harvest interval on edible crops.

Please consult MSDS and product label prior to use.

Hortifeeds Direct, Park Farm, Kettlethorpe, Lincoln.

Tel. 01522 704404 Fax. 01522 704748 email, info@hortifeeds.co.uk www.hortifeeds.co.uk